

An energetic stellar outburst accompanied by circumstellar light echoes

Howard E. Bond¹, Arne Henden², Zoltan G. Levay¹ Nino Panagia^{1,3},
William B. Sparks¹, Sumner Starrfield⁴, R. Mark Wagner⁵, R. L. M. Corradi⁶,
& U. Munari⁷

Received _____; accepted _____

To appear in Nature, March 27, 2003

¹Space Telescope Science Institute, 3700 San Martin Dr., Baltimore, MD 21218, USA

²Universities Space Research Association & US Naval Observatory, Flagstaff Station, P.O. Box 1149, Flagstaff, AZ 86002, USA

³On assignment from European Space Agency, Space Telescope Division

⁴Dept. of Physics & Astronomy, Arizona State University, Tempe, AZ 85287-1504, USA

⁵Large Binocular Telescope Observatory, University of Arizona, 933 North Cherry Ave., Tucson, AZ 85721, USA

⁶Isaac Newton Group of Telescopes, Apartado de Correos 321, 38700 Santa Cruz de La Palma, Canarias, Spain

⁷INAF-Osservatorio Astronomico di Padova, Sede di Asiago, 36012 Asiago (VI), Italy

Some classes of stars, including supernovae and novae, undergo explosive outbursts that eject stellar material into space. In 2002, the previously unknown variable star V838 Monocerotis brightened suddenly by a factor of $\sim 10^4$. Unlike a supernova or nova, V838 Mon did not explosively eject its outer layers; rather, it simply expanded to become a cool supergiant with a moderate-velocity stellar wind. Superluminal light echoes were discovered^{1,2} as light from the outburst propagated into surrounding, pre-existing circumstellar dust. Here we report high-resolution imaging and polarimetry of the light echoes, which allow us to set direct geometric distance limits to the object. At a distance of >6 kpc, V838 Mon at its maximum brightness was temporarily the brightest star in the Milky Way. The presence of the circumstellar dust implies that previous eruptions have occurred, and spectra show it to be a binary system. When combined with the high luminosity and unusual outburst behavior, these characteristics indicate that V838 Mon represents a hitherto unknown type of stellar outburst, for which we have no completely satisfactory physical explanation.

Light echoes around an outbursting star appear after the eruption, the illumination being delayed by the longer path length from the star to the surrounding dust and then to the Earth, as compared with the light from the outburst itself traveling directly to the Earth. In the case of an instantaneous light pulse, the geometry of a light echo is simple^{3–8}: the illuminated dust lies on the paraboloid given by $z = x^2/2ct - ct/2$, where x is the projected distance from the star in the plane of the sky, z is the distance from this plane along the line of sight toward the Earth, c is the speed of light, and t is the time since the outburst. Light echoes are rare, having been observed only around a few classical novae in our own Galaxy^{3,9,10}, and around several extragalactic supernovae¹¹. With the exception of SN 1987A¹², these echoes arose from interstellar dust or gas in the line of sight, rather than

from true circumstellar material.

V838 Mon erupted in early 2002 January¹³. Fig. 1 shows the light curve of the outburst in linear flux units, from ground-based observations in three different wavelength bands: blue (B), visual (V), and near-infrared (I). This light curve—with a weak precursor brightening, a sharp, bright blue peak, followed by a decline and then two further broad and much redder peaks—is unlike that of a supernova, nova, or any other known type of variable star. Spectra reported in numerous *IAU Circulars* showed a very cool photosphere throughout the outburst. Toward the end of the outburst, and during the decline, the star became extremely red. Recent spectra^{20–22}, taken after the decline, show that a hot B-type star is also present, along with the cool outbursting object.

We have obtained *Hubble Space Telescope* (*HST*) observations of the light echoes surrounding V838 Mon with the Advanced Camera for Surveys (ACS) at the four epochs listed in Table 1. Fig. 2 shows the ACS images. Only the B filter was used at the first epoch, but for the others we obtained three-color (BVI) frames, and have used these to make renditions that approximate true color. The nebulosity exhibits a wealth of subarcsecond cirrus-like structure, but is dominated (especially at the first two epochs) by a remarkable series of nearly circular arcs and rings, centered on the variable star. The exact linear size of the illuminated dust envelope depends on the distance to the object, but based on the distance limits given below, the visible material is confined to less than 2 pc from the star and is thus circumstellar, rather than being ambient interstellar medium. The cavity centered on the star, e.g. in Figs. 2a and 2b, along with several features that appear aligned toward the star, also establish the circumstellar nature of the echoes.

As shown in Fig. 1, the outburst light curve had the simplest structure in B —a bright, relatively short peak on February 6, and a second, weaker peak about 32 days later—making the B -band light echoes most amenable to geometric analysis. If we assume

the simplest geometry, which is a series of nested spherical dust shells centered on the star and intersecting the expanding light paraboloids, then it can be shown that the apparent expansion rate of a ring is given by $dx/dt = (rc - c^2t)(2rct - c^2t^2)^{-1/2}$, where r is the radius of the ring measured from the star. Thus the time behavior of these rings in angular units *provides a direct geometric distance to the object*. The pair of images taken only 20.8 days apart, on April 30 and May 20, is the most suitable for such a measurement, since the same rings can be identified unambiguously in both images. The expansion rates are plotted against the mean angular radii of each ring in Fig. 3. Overlaid on the plot are the expansion rates calculated from the above equation, for ages of 62 and 93 days, which are the elapsed times between the first and second B -band light-curve peaks and the mean epoch of the two images. It can be seen that the measured points fall into two families, plotted for clarity as filled and open circles. The filled circles satisfactorily follow the expected behavior if the corresponding echo rings are illuminated by the first light-curve peak, while the open circles appear to represent rings illuminated by the second peak. In strong confirmation of this interpretation, we find that the filled circles all refer to the blue rings seen clearly in Fig. 2b, while the open circles all correspond to the trailing, red-edged rings.

Fig. 3 does show significant scatter, probably arising from departures of the actual circumstellar structures from the simple adopted spherical geometry. For plausible non-spherical geometries, measures made over several azimuths will still average to a result close to the true distance. Thus, barring a very unusual morphology, the figure implies that the distance to V838 Mon must certainly be greater than 2 kpc, and probably significantly greater. Since the predicted expansion rates approach an asymptotic limit with increasing distance, we cannot make a stronger statement at present. However, because the relations for various distances diverge with increasing time, continued high-resolution imaging will eventually yield a very precise distance.

A second, independent geometrical method for determining the distance comes from polarimetry⁸. Since maximum linear polarization occurs for 90° scattering, there will be highly polarized light at a linear radius ct , whose angular radius would thus yield the distance—but of course only if there actually is dust at that radius and at the same distance from us as the star. Our most recent polarimetric *HST* image, obtained on 2002 September 2, shows very high polarization ($\sim 50\%$), but only at the inner rim of the cavity at the lower left (southeast) of the star. Thus, at present, we can only set a lower limit to the distance, by equating the radius at this rim, $\sim 5''.0$, to ct , where t is now the time since the second light-curve peak. This lower limit is ~ 6 kpc. Future observations will provide a definitive distance from this method as well, once the light-echo paraboloid enlarges enough to allow us to illuminate close-in material behind as well as in front of the star. At that time, we expect to see maximum polarization in a ring⁸, rather than right at the inner edge of the rim.

For a distance of at least 6 kpc (and interstellar reddening of $E(B - V) \simeq 0.8^{16,23}$), the absolute magnitude of V838 Mon at maximum was at least $M_V = -9.6$, making it more luminous than a classical nova, and temporarily the brightest star in the Milky Way. The light curve, spectroscopic behavior, and high luminosity of V838 Mon are reminiscent of “M31 RV,” a luminous red variable star that appeared in the Andromeda galaxy in the late 1980’s²⁴. M31 RV reached an absolute visual magnitude at least as bright as $M_V = -9.3^{25}$, but unfortunately was not well studied before it declined below detectability. V838 Mon and M31 RV (along with, possibly, the Galactic object V4332 Sagittarii²⁶) appear to represent a new type of outburst in which a star expands rapidly to supergiant dimensions²⁷, but without either the catastrophic envelope loss or the evolution to high temperature seen in classical and symbiotic novae.

In the case of V838 Mon the outburst appears to have occurred in a binary system,

which may suggest an analogy with a nova system in which a thermonuclear explosion occurs on an accreting white-dwarf companion. However, a broad theoretical exploration²⁸ of the parameter space for thermonuclear runaways on accreting white dwarfs predicts no behavior like that of V838 Mon. Evidently, V838 Mon has in the recent past had several similar outbursts, which produced the dust envelope that is now being illuminated. These previous outbursts appear to rule out one-time catastrophic events such as a stellar collision or merger²⁹, or a common-envelope interaction³⁰, as the energy source. Now that V838 Mon appears to be returning to its quiescent state, further observations of the system may shed light on its properties and give insight into possible outburst mechanisms for this mysterious and extraordinary object.

References:

1. Henden, A. et al. V838 Monocerotis. *IAU Circular* 7859 (2002).
2. Munari, U. et al. The mysterious eruption of V838 Mon. *Astron. Astrophys.* **389**, L51-L56 (2002).
3. Schaefer, B.E. Light echoes: novae. *Astrophys. J.* **327**, 347-349 (1988).
4. Kapteyn, J.C. Über die Bewegung der Nebel in der Umgebung von Nova Persei. *Astr. Nachrichten* **157**, 201-204 (1902).
5. Couderc, P. Les auréoles lumineuses des novae. *Ann. d'Astrophys* **2**, 271-302 (1939).
6. Russell, H.N., Dugan, R.S. & Stewart, J.Q. *Astronomy: a revision of Young's Manual of Astronomy* (Ginn & Co., Boston, 1945), p. 786.
7. Chevalier, R.A. The scattered-light echo of a supernova. *Astrophys. J.* **308**, 225-231 (1986).
8. Sparks, W.B. A direct way to measure the distances of galaxies. *Astrophys. J.* **433**, 19-28 (1994).
9. Ritchey, G.W. Nebulosity about Nova Persei. Recent photographs. *Astrophys. J.* **15**, 129-131 (1902).
10. Casalegno, R. et al. The emission nebula associated with V1974 Cygni: a unique object? *Astron. Astrophys.* **361**, 725-733 (2000).
11. Sparks, W.B. et al. Evolution of the light echo of SN 1991T. *Astrophys. J.* **523**, 585-592 (1999).
12. Bond, H.E., Gilmozzi, R., Meakes, M.G. & Panagia, N. Discovery of an inner light-echo ring around SN 1987A. *Astrophys. J.* **354**, L49-L52 (1990).
13. Brown, N.J. Peculiar Variable in Monoceros. *IAU Circular* 7785 (2002).
14. Kato, T. <http://vsnet.kusastro.kyoto-u.ac.jp/vsnet/index.html> (2002).
15. Goranskii, V.P. et al. Nova Monocerotis 2002 (V838 Mon) at early outburst stages. *Astronomy Letters* **28**, 691-700 (2002).

16. Kimeswenger, S., Lederle, C., Schmeja, S. & Armsdorfer, B. The peculiar variable V838 Mon. *Monthly Notices Roy. Astron. Soc.* **336**, L43-L47 (2002).
17. Osiwala, J.P. et al. The double outburst of the unique object V838 Mon. In *Symbiotic stars probing stellar evolution*, eds. R.L.M. Corradi, J. Mikolajewska & T.J. Mahoney eds., (ASP Conference Series, San Francisco, 2002), in press.
18. Price, A. et al. Multicolor observations of V838 Mon. *IAU Inform. Bull. Var. Stars* 5315, 1 (2002).
19. Crause, L.A., Lawson, W.A., Kilkenny, D., van Wyk, F., Marang, F. & Jones, A.F. The post-outburst photometric behaviour of V838 Mon, *Monthly Notices Roy. Astron. Soc.* in press.
20. Geballe, T.R., Smalley, B., Evans A. & Rushton, M.T. V838 Monocerotis. *IAU Circular* 8016 (2002).
21. Munari, U. & Desidera, S. V838 Monocerotis. *IAU Circular* 8005 (2002).
22. Wagner, R.M. & Starrfield, S. V838 Monocerotis. *IAU Circular* 7992 (2002).
23. Zwitter, T. & Munari, U. V838 Monocerotis. *IAU Circular* 7812 (2002).
24. Mould, J. et al. A nova-like red variable in M31. *Astrophys. J.* **353**, L35-L37 (1990).
25. Bryan, J. & Royer, R.E. Photometry of the unique luminous red variable in M31. *Publ. Astron. Soc. Pacific* **104**, 179-181 (1992).
26. Martini, P. et al. Nova Sagittarii 1994 1 (V4332 Sagittarii): The discovery and evolution of an unusual luminous red variable star. *Astron. J.* **118**, 1034-1042 (1999).
27. Munari, U., Henden, A., Corradi, R.L.M. & Zwitter, T. V838 Mon and the new class of stars erupting into cool supergiants. In *Classical Nova Explosions*, eds. M.Hernanz & J.Jose (American Institute of Physics, New York, 2002), p. 52-56.
28. Prialnik, D. & Kovetz, A. An extended grid of multicycle nova evolution models. *Astrophys. J.* **445**, 789-810 (1995).

29. Soker, N. & Tylenda, R. Main-sequence stellar eruption model for V838 Monocerotis. *Astrophys. J.* **582**, L105-L108 (2003).
30. Iben, I. & Tutukov, A.V. Rare thermonuclear explosions in short-period cataclysmic variables, with possible application to the nova-like red variable in the Galaxy M31. *Astrophys. J.* **389**, 369-374 (1992).

Acknowledgements Based on observations with the NASA/ESA *Hubble Space Telescope*, obtained at the Space Telescope Science Institute, which is operated by AURA, Inc., under NASA contract NAS5-26555. We thank the Space Telescope Science Institute for awarding Director’s Discretionary Observing Time for this project and for support through grant GO-9587. S.S. was supported, in part, by NSF and NASA grants to Arizona State University.

Competing interests statement The authors declare that they have no competing financial interests.

Correspondence and requests for materials should be addressed to H.E.B. (e-mail: bond@stsci.edu).

Table 1. *Hubble Space Telescope* ACS Wide-Field Channel Observing Log

UT Date (2002)	HJD	Days since Feb 6.4	Days since Mar 10.0	ACS filters used
Apr 30	2452394.7	82.8	51.2	B +pol
May 20	2452415.5	103.6	72.0	B +pol, V +pol, I
Sep 2	2452520.4	208.5	176.9	B , V +pol, I
Oct 28	2452576.0	264.1	232.5	B , V , I

The Advanced Camera for Surveys’ F435W, F606W, and F814W filters are denoted B , V , and I , respectively, and “+pol” denotes that the indicated filter was combined in succession with 0-, 60-, and 120-degree polarizers. The time of maximum B luminosity was near 2002 February 6.4, and the second B peak was near March 10.0. (The lack of *HST* observations between May and September is due to solar avoidance.)

FIGURE CAPTIONS

Fig. 1.— Multicolor light curves of the outburst of V838 Mon. Data have been assembled by A.H. from observations by himself at the U.S. Naval Observatory, Flagstaff, and from amateurs¹⁴ and other professional observers^{2,15–19}. Magnitudes in the standard Johnson blue (B), visual (V), and Kron-Cousins near-infrared (I) bands have been converted to flux on a linear scale, normalized to the peak at Heliocentric Julian Date 2452311. Corresponding calendar dates are shown at the bottom. After an initial rapid rise to 10th magnitude and a subsequent gradual decline, V838 Mon abruptly rose another 4 magnitudes in early February, reaching a maximum of visual magnitude $V = 6.75$ on 2002 February 6. After another decline, the star rose to a second peak in early March. Once again there was a slow decline after this peak, except in the near-infrared I band, where yet a third peak was attained in early April. In late April, V838 Mon dropped quickly back to its original quiescent brightness in the blue (B) and V bands; this precipitous drop in brightness has been attributed to dust formation above the photosphere¹⁹. The most recent observations show some rebrightening in I only. The progenitor object was recorded in archival sky surveys² at visual magnitude 15.6, very similar to its current post-outburst brightness.

Fig. 2.— *Hubble Space Telescope* Advanced Camera for Surveys images of the light echoes. The apparently superluminal expansion of the echoes as light from the outburst propagates outward into surrounding dust is shown dramatically. Images were taken on (a) 2002 April 30, (b) May 20, (c) September 2, and (d) October 28. Each frame is $83'' \times 83''$; north is up and east to the left. Imaging on April 30 was obtained only in the B filter, but B , V , and I were used on the other three dates, allowing us to make full-color renditions. The time evolution of the stellar outburst (Fig. 1) is dramatically reflected by structures visible in the color images. In Fig. 2b, for example, note the series of rings and filamentary structures, especially in the upper right quadrant. Close examination shows that each set of rings has a sharp, blue outer edge, a dip in intensity as one moves toward the star, and then a rebrightening to a redder plateau. Similar perfect replicas of the outburst light curve are seen propagating outwards throughout all of the color images.

Fig. 3.— Measured and predicted apparent angular expansion rates of rings seen in the light echoes. Measurements were done on the pair of *B*-band images taken on 2002 Apr 30 and May 20. A simple edge-finding algorithm was used to mark the locations of the same rings on both images at several different azimuths, and their angular expansion rates were calculated. Filled circles denote rings that we attribute to illumination from the first light-curve peak, and open circles denote rings attributed to the second peak. Measurement errors are only slightly larger than the size of the plotted points. Units of expansion are milliarcseconds per day, as functions of angular radius in arcseconds. Also shown are the predicted expansion rates for light-echo rings seen 62 days (red dashed lines) and 93 days (blue solid lines) after outburst, calculated from the equation given in the text. They are functions of distance to the star, plotted for the range from 0.5 to 16 kpc, and are apparently superluminal due to the parabolic geometry of the illuminated surfaces. The measured angular expansion rates imply a strong lower limit to the stellar distance of 2 kpc.




Figure 1




Figure 2




Figure 3